



News Blogs

Email Login Log In

Forgot password? Not a member?

All Health Bioscience Physical Science Environment Space Tech Life Origins Misc

Press Release

Blood progenitor cells receive signals from niche cells and the daughter blood cells they create

Thursday, December 22, 2011

Maintaining balance is crucial. In *Drosophila*, the common fruit fly, the creation and maintenance of the blood supply requires such balance.

UCLA stem cell scientists have now uncovered that two-way signaling from two different sets of cells is necessary for that balance, both to ensure enough blood cells are made to respond to injury and infection and that the blood progenitor cell population remains available for future needs.

The stem cell-like blood progenitor cells – which contribute to the cells of the adult fruit fly's blood supply – receive signals from cells that live in a nearby safe zone, or niche. These signals keep the progenitors in the same stem cell-like state so, when needed, they can begin differentiating into blood cells.


And in a new discovery, the UCLA stem cell scientists found that the blood progenitor cells receive critical signals back from the daughter blood cells they create, telling the progenitor cells when enough blood cells have been made and it's time to stop differentiating.

The new discovery of the "back talk" from the daughter blood cells appears Dec. 23, 2011 in the peer-reviewed journal *Cell*.

"The cells in the niche provide a safe environment to support blood progenitor cells," said study co-senior author Dr. Julian A. Martinez-Agosto, an assistant professor of human genetics and pediatrics and a researcher with the Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research at UCLA. "When the blood progenitor cells receive signals from the niche cells it creates an environment for those cells to maintain their potential and not differentiate."

Previous studies have shown that when you remove the niche cells, the blood progenitor cells differentiate unchecked. Ultimately, the fruit fly runs out of blood progenitor cells and is not able to make new blood cells to mount an immune response to infection or injury, Martinez-Agosto said.

The new findings by Martinez-Agosto and study co-senior author Utpal Banerjee, a Broad center researcher and the Irving and Jean Stone Professor and chairman of molecular, cell and developmental biology in Life Sciences, identified additional signals not coming from the niche cells. The new signals were coming from the daughter blood cells the progenitors were making,



Red blood cells

Labspaces.net on Facebook

Like

1,590 people like **Labspaces.net**.

Husam 妍彤 Rachel Paul David

Facebook social plugin

Featured Blogs

Got fetal DNA on the brain?



Author: **Brian Krueger, PhD** | Comments: 3

Sight without seeing: Balint's syndrome



Author: **Jordan Gaines** | Comments: 0

Memes about 'junk DNA' miss the mark on paradigm shifting science



Author: **Brian Krueger, PhD** | Comments: 2

Catnip fever: why your cat acts high



Author: **Jordan Gaines** | Comments: 2

So, you've dropped a vial or lost a sample box in your liquid nitrogen container...now what?



Author: **Brian Krueger, PhD** | Comments: 2

[View more Blog articles](#)

More Biological Science

The genetics of white finger disease



genitor cells receive signals from niche cells and the daughter blood cells they create

a surprising discovery, Banerjee said.

Martinez-Agosto and Banerjee noted in the four-year study that once the progenitors cells had begun differentiating and the blood cells they were creating became mature, the progenitors became very quiescent, or quiet, and did not multiply. They theorized that there must be a signal coming from the daughter cells that told the progenitors to stop multiplying and differentiating.

"It was a very surprising finding, because there was no reason to suspect that the differentiating cells had any role at all in the process," Banerjee said. "It's always been the paradigm in stem cell biology that all that was needed was the signaling from the niche cells to maintain the progenitor population. Now, we've shown that you also need the signals from the daughter cells to help maintain the progenitor cell population."

The signaling from the niche cells that maintains the progenitor population is called Hedgehog. In this study, the scientists showed that the daughter cells are sending back a signal to the progenitors that is mediated by Adenosine deaminase growth factor A (Adgf-A). The signal regulates extracellular levels of adenosine, which opposes or counters the effects of Hedgehog signaling.

"We've shown that adenosine as a molecule is really important for regulating the proliferation of progenitor cells in blood. And it requires a delicate balance – just enough signaling to give you more blood cells, but not so much that all the progenitor cells are lost," Martinez-Agosto said. "Maybe other progenitors or stem cells are using the same signaling to determine when to differentiate or not."

The team used the fruit fly because it is a very accessible model organism in which genes can be easily manipulated and their effects on cells monitored, Martinez-Agosto said. They dissected the fly lymph gland, where blood cells are made, and used green fluorescence to label progenitors and their daughter cells to determine when they were differentiating.

Going forward, the team will try to understand if the progenitor cells can sense the adenosine in their microenvironment under stress and injury conditions and how cell division biologically counters the niche signaling to promote formation of blood cells.

The study was funded in part by the National Heart, Lung and Blood Institute.

"Our findings reveal signals arising from differentiating cells that are required for maintaining progenitor cell quiescence and that function with the niche-derived signal in maintaining the progenitor state," the study states. "Similar homeostatic mechanisms are likely to be utilized in other systems that maintain relatively large numbers of progenitors that are not all in direct contact with the cells of the niche."

###

University of California - Los Angeles Health Sciences: <http://www.uclahealth.org/>

Thanks to University of California - Los Angeles Health Sciences for this article.

This press release was posted to serve as a topic for discussion. Please comment below. We try our best to only post press releases that are associated with peer reviewed scientific literature. Critical discussions of the research are appreciated. If you need help finding a link to the original article, please contact us on twitter or via e-mail.

This press release has been viewed 106 time(s).



0

submit
reddit


2
Tweet

0
Share

Comments

Source: BioMed Central | Comments: 0

Scientists find missing link between players in the epigenetic code



Source: University of North Carolina Health Care | Comments: 0

Common RNA pathway found in ALS and dementia



Source: University of California - San Diego | Comments: 0

Researchers discover gene that causes deafness



Source: University of Cincinnati Academic Health Center | Comments: 0

Marine plants can flee to avoid predators




Source: University of Rhode Island | Comments: 0

White shark diets vary with age and among individuals




Source: University of California - Santa Cruz | Comments: 0

Researchers find multiple similarities between cancer cells and induced pluripotent stem cells



Source: University of California - Davis Health System | Comments: 0

Penn immunologists find a molecule that puts the brakes on inflammation



Source: University of Pennsylvania | Comments: 0

From the Web

Great white sharks off West Coast to be considered for endangered listing

The federal government will examine whether to protect the West Coast population of great white sharks under the Endangered Species Act, the National Oceanic and Atmospheric Administration announced Thursday.

[Marine Biology](#) | Source: Washington Post | Views: 59 | Comments: 0

Burn victim identified by maggots on body

Police have discovered the identity of a burned body using DNA extracted from the guts of maggots feeding on the victim

[Genetics](#) | Source: New Scientist | Views: 50 | Comments: 0

Forget the Cloud—Knome Offers Genome Analysis in a Box

A \$125,000 hardware-and-software system could help researchers analyze data securely and with less IT support.

[Genetics](#) | Source: Technology Review | Views: 47 | Comments: 0

New species of plankton discovered

Microscopic creatures at the bottom of the food chain play an over-sized role in the global ecosystem

[Marine Biology](#) | Source: CBSNews | Views: 47 | Comments: 0

ScienceShot: To Feast on Toxic Plant, Insects Find Common Genetic Ground

Distantly related species took parallel genetic strategies to survive milkweed

[Evolution](#) | Source: Science | Views: 54 | Comments: 0

N

Add Comment?

Path: p

For youtube videos, paste embed code directly in the text box

- Notify me by email when future posts are made

Email:

Members do not need to provide an address

- Preview before posting

Select Comment Validation Method

- Member
- Name/URL (Guest)
- FaceBook (Guest)

Rate Article

Total votes: 0

Recognition at last for Alfred Russel Wallace, who lived in Darwin's shadow

Victorian naturalist's extensive collection of specimens and papers appear together for first time in online project

Evolution | Source: TheGuardian | Views: 72 | Comments: 0

Taxidermists, Artists Bring Aging Museum Animals Back to Lifelike

The North American mammals at the American Museum of Natural History are coming back to (near) life. Artists, taxidermists, conservators and designers have been working for a year on everything from re-coloring faded bear fur and restoring paintings of the Grand Canyon to dusting leaves and installing new energy-efficient and less-damaging lighting.

Animals | Source: Wired | Views: 28 | Comments: 0

Flying backwards is efficient

Flying backwards uses surprisingly little energy, according to a recent scientific study.

Animal Behavior | Source: BBC News | Views: 58 | Comments: 0

Tiny seahorse spotted in underwater Google Street View

This pygmy seahorse was confirmed in Australian waters for the first time during the underwater Catlin Seaview Survey, which is mapping the world's reefs

Marine Biology | Source: New Scientist | Views: 36 | Comments: 0

The Great Barrier Reef is heating up

The waters of Australia's Great Barrier Reef are getting warmer, satellite measurements show, which spells bad news for the myriad creatures that dwell in this rain forest of the ocean.

Marine Biology | Source: NBCnews | Views: 22 | Comments: 0

Friends



Facebook



Carver College of Medicine



Twitter



LabX.com



LabManager.com



ADVANCE Magazine



BioData.com



Adamson Studios



Drew Curtis' Fark



MicrobeWorld



NYAS



Iowa City Web Services