

[Sign in](#)[Register](#)

- [Phys.org](#)
- - [Main page](#)
  - [Feature stories](#)
  - [Weblog & Reports](#)
  - [Archive](#)
  - [Video](#)
  - [Podcasts](#)
- [Follow us](#)
  - [Facebook](#)
  - [Twitter](#)
  - [RSS feeds](#)
- [Mobile Apps](#)
  - [iPhone](#)
  - [Android](#)
    - [Medical Xpress \(free\)](#)
    - [Medical Xpress](#)
  - [BlackBerry](#)
  - [Amazon kindle](#)
- [Search](#)
  - 
  - [advanced search](#)

[Medical, health and wellness news](#)

- [Home](#)
- [Topics](#)
- [Conditions](#)
- [Latest news](#)
- [Week's top](#)
- [Other news](#)
- [Spotlight news](#)
- [News w/ video](#)
- [Podcasts](#)
- [Unread news](#)
- [Meet Medical Xpress: PhysOrg.com transfers medical section into a separate site. \[Read more\]\(#\)](#)

## Tumor suppressor genes vital to regulating blood precursor cells in fruit flies

September 5, 2012 in [Medical research](#)

**UCLA stem cell scientists have shown that two common tumor suppressor genes, TSC and PTEN, are vital to regulating the stem cell-like precursor cells that create the blood supply in *Drosophila*, the common fruit fly.**

[Ads by Google](#)

[Nothing but the web.](#) - Chromebooks are built and optimized for the web. Learn more. - [google.com/chromebook](http://google.com/chromebook)

The researchers examined a signaling pathway called TOR that the cells use to gauge nutrition levels and stress, said study senior author Dr. Julian A. Martinez-Agosto, an assistant professor of [human genetics](#) and pediatrics and a researcher with the Eli and Edythe Broad Center of Regenerative Medicine and [Stem Cell Research](#) at UCLA.

"We wondered how an organism knows how many [blood cells](#) to make and when to make them in the context of injury and repair to tissue. In particular, we wondered how the blood progenitor cells sense that change and know when it's time to make more blood cells," Martinez-Agosto said. "We found that the TOR pathway uses these two genes to regulate its function and, when activated, it expands or increases the number of blood progenitor cells in the fly's blood."

The study appears Sept. 5, 2012 in the advance online issue of the peer-reviewed journal *Development*.

Michelle Dragojlovic-Munther, a graduate student in the Martinez-Agosto laboratory and first author of the study, found that cells with increased activity of TOR have a competitive advantage, allowing them to divide and make more of themselves so they can make blood. These progenitors, Dragojlovic-Munther found, also have high levels of reactive [oxygen species](#) (ROS) - ions or very small molecules that include [free radicals](#) - which are known to [damage cells](#) and can predispose humans to aging and heart disease. But in this case, the ROS proved valuable.

The precursors, Martinez-Agosto said, were producing ROS all the time and when TOR was activated, the levels increased dramatically. Too much ROS caused them to divide more than normal. If they treated the flies with antioxidants, which reduce ROS levels, the cells would develop normally.

The finding could be important because the TOR pathway is abnormally activated in many cancers, and it may be possible to target the levels of ROS, which may help regulate the pathway.

"What this study may be telling us is that too much ROS is causing more cells to divide and we may be able to target therapies that reduce ROS to significantly improve the condition," Martinez-Agosto said, adding that specifically targeted antioxidants might be a potential treatment in certain subsets of blood disorders. "Sometimes that pathway is working more

than it should, and we need the right amount of ROS for balance. It's like Goldilocks, there can't be too little or too much. We need it just right."

Going forward, Martinez-Agosto and his team will try to determine where the ROS is coming from and perhaps discover an enzyme that may be a good target for therapeutics. They know that the higher ROS levels in blood progenitors are not coming from mitochondria, the cell's power source, but have not identified how they are being produced.

"This study highlights mechanistic differences between TSC and [PTEN](#) on TOR function and demonstrates the multifaceted roles of a nutrient-sensing pathway in orchestrating proliferation and differentiation of myeloid-specific blood [progenitors](#) through regulation of ROS levels and the resulting myeloproliferative disorder when deregulated," the study states.

**Journal reference:** [Development](#)

**Provided by** [University of California, Los Angeles](#)



1 simple rule to making a fortune overnight...



Advertisement  
Scientists in Cambridge have discovered a revolutionary new muscle builder...



The secret of how to learn a foreign language in just 10 days. Read here to find out...



Men over 40: Feeling flabby, weak, and struggling with low libido? Take back your life with this.

[send feedback to editors](#)

not rated yet

- [1](#)
- [2](#)
- [3](#)
- [4](#)
- [5](#)

Please [register](#) or [sign in](#) to add a comment. Registration is free, and takes less than a minute. [Read more](#)

Sign in with

Email

Password

Sign In

Forgot your password? [Click here](#) to reset it.

Notify me via email of follow-up comments posted here  [sign in first](#)

Rank

- [1](#)
- [2](#)
- [3](#)